



# Loudon Chapter 17 Review: Allylic/Benzylic Reactivity

Jacque Richardson, CU Boulder – Last updated 9/19/2016


Chapter 17 is all about reactions that happen at the position one away from an aromatic ring, or one away from a double bond. These are called the benzylic and allylic positions respectively. Benzyl and allyl are the names of the corresponding R groups.



The most important thing about both these positions is that they're great at stabilizing any sort of charge or radical, through resonance. The allyl group has two resonance forms:





The benzyl group has four resonance forms (or possibly more, if there are any functional groups attached to the ring):



What this means is that any reaction that involves a charge or a partial charge forming at the allylic or benzylic position happens much more easily than it would for a regular R group. This includes  $\text{S}_{\text{N}}1$ ,  $\text{S}_{\text{N}}2$ , E1, and E2, along with other reactions that we've seen in the past.

## Reactions with Carbocations


The biggest two in this category are  $\text{S}_{\text{N}}1$  and E1. Here are the mechanisms for benzylic only, but they look similar for allylic. None of the resonance forms are shown for the benzylic cation but they are responsible for its stability.




In both cases, the reaction goes much faster because the high-energy intermediate is stabilized. The reaction is usually a few hundred times faster than it would be on a saturated alkyl halide. This effect can become even greater if there are groups on the ring that stabilize the positive charge.

# Loudon Chapter 17 Review: Allylic/Benzylic Reactivity


Jacque Richardson, CU Boulder – Last updated 9/19/2016



The intermediates here look somewhat similar to the intermediates for EAS reactions in Ch. 16. Since we're forming a carbocation, just like in EAS, it's not surprising that the same rules will apply for activating/deactivating groups. Also, a group will have a much stronger effect if it's ortho or para to where the benzylic carbon is attached. In the example above, the hydroxyl (OH) group is para to the benzylic carbon and so it's capable of putting the charge directly onto the oxygen. To rank some examples by how fast they react at  $S_N1/E1$ :




One more complication of resonance stabilization: if you make a carbocation at the allylic position, you can theoretically get two different products. This is a lot like the 1,2 vs. 1,4 additions from Ch. 15 – at some point, there are two different resonance forms to react with. Which one gives the major product depends on where the charge is most stable – here, the second product is the major one because the tertiary carbocation is the major resonance form.



This is not the case, however, for reactions at the benzylic position. There's only one resonance form that's actually aromatic, in those cases, so that's the only one that leads to product.

## Reactions with Radicals

The allylic and benzylic positions stabilize radicals just as well as carbocations. We saw radical halogenation in Ch. 9, which uses  $Br_2/h\nu$  to replace an H with a Br. This happens almost exclusively at the carbon that can best stabilize a radical. In alkanes this was only on the most substituted carbon (usually  $3^\circ$ ), but the allylic and benzylic positions are even better.



In this case, even though there's a  $3^\circ$  carbon at the right end of the molecule, the benzylic position is still preferred. There's a problem if we try to do this reaction on the allylic position, though.  $Br_2$  can add straight to the alkene if it doesn't go by a radical mechanism.

# Loudon Chapter 17 Review: Allylic/Benzylic Reactivity


Jacque Richardson, CU Boulder – Last updated 9/19/2016



If it goes by radical mechanism

If it ignores  $h\nu$  and goes by regular addition mechanism


Usually we can favor the radical mechanism by using a lot of light, heat, and a polar solvent, and we can favor the addition mechanism by the opposite conditions. There are some problems in the book that say “Br<sub>2</sub>/dark”, for instance, avoiding radical formation. There’s a new reagent that shows up here: N-bromosuccinimide, or NBS. (There are also NCS and NIS, for chlorine and iodine respectively, though NBS works best.)




It works as a source of Br radicals without ever adding two Br atoms to the molecule the way that Br<sub>2</sub> would. Also unlike Br<sub>2</sub>, it goes only for the benzylic/allylic position, and doesn’t react significantly with tertiary carbons. Again, if we do this reaction at the allylic position, a mixture of products is possible based on different resonance forms.

## Reactions with Carbanions

Again, a negative charge at the allylic or benzylic position is much more stable than it would be in a lot of other positions. We’ve seen carbanions before in the form of Grignards or organolithiums. In those cases, we could show the carbon-metal bond as either covalent or as an ionic pair – both are acceptable. Here, the charge can move around to become what looks like a different structure of organometallics, though it’s all just different depictions of the same thing.




The molecule can behave as if it were either one of these structures, so if we react it with something that takes away its charge and resonance it can be trapped in either form.



## Reactions without Formal Charges

Even if there’s no official positive or negative charge, reactions at this position can still have stabilized partial charges. S<sub>N</sub>2 and E2 fit this description. In the reaction below, you’d expect, given the combination of strong base/good nucleophile/primary alkyl halide, to see S<sub>N</sub>2 as the major pathway. But it turns out to be E2 instead.




# Loudon Chapter 17 Review: Allylic/Benzylic Reactivity

Jacque Richardson, CU Boulder – Last updated 9/19/2016


This is because E2 creates a slight negative charge at the benzylic position, which stabilizes the reaction quite a lot and allows it to happen faster. If the mechanism were S<sub>N</sub>2, the benzylic position wouldn't be involved at all.

If the halogen itself is located at the benzylic/allylic position instead of one carbon away from it, though, then S<sub>N</sub>2 becomes extremely favored. For allylic, it gets sped up by a couple of orders of magnitude, but for benzylic it gets sped up by about five orders of magnitude. This has less to do with charge stabilization and more to do with stabilization of the sp<sup>2</sup>-hybridized intermediate. The p orbitals of the atom doing the reaction can overlap with the p orbitals of the double bond/aromatic ring.




## Allylic & Benzylic Oxidations

We're used to seeing alcohols getting oxidized by PCC or strong oxidizers like CrO<sub>3</sub>/H<sub>2</sub>SO<sub>4</sub>:




These will always do predictable things to an alcohol depending on whether it's 1°, 2°, or 3°, regardless of what the attached R groups actually are. In this chapter we start looking at reagents that specifically target alcohols in the allylic or benzylic positions, without touching other alcohol groups.



Note that if we start with a primary allylic/benzylic alcohol, it still maxes out at two bonds to oxygen (the aldehyde/ketone oxidation level) without going all the way to three bonds to oxygen (the carboxylic acid oxidation level). For this reason, we can think of MnO<sub>2</sub> as being kind of like a version of PCC that only goes after allylic/benzylic alcohols.

Another oxidation reaction that specifically targets these locations doesn't start with an alcohol at all. You can use either KMnO<sub>4</sub>, CrO<sub>3</sub>, or Na<sub>2</sub>Cr<sub>2</sub>O<sub>7</sub>, but they need vigorous conditions – heat and strong acid – to work.



## Loudon Chapter 17 Review: Allylic/Benzylic Reactivity

Jacque Richardson, CU Boulder – Last updated 9/19/2016

The general idea here is that if we have a carbon attached to an aromatic ring, and that carbon has at least one hydrogen or OH group on it, it will be oxidized all the way to a carboxylic acid – *even if that means losing bonds to carbons that it's attached to*. This is not something that we've seen oxidizers do before. They've always left carbon-carbon bonds intact until now, but  $\text{KMnO}_4$  is a very powerful oxidizer and the benzylic position is pretty reactive.